

ZENBEXUS™ REMS | Pharmacy Counseling Checklist

For Females Who Can Become Pregnant and Males

All pharmacies certified by ZENBEXUS REMS (Risk Evaluation and Mitigation Strategy) must complete this form prior to each dispense of ZENBEXUS™ (iberdomide) for females who can become pregnant and males.

Submit this completed checklist via:

- **Online** at www.BMSREMSPatientSafety.com **or**
- **Email** to REMSContactCenter@bms.com **or**
- **Fax** to 1-877-296-1932

Do not dispense ZENBEXUS until the REMS provides confirmation to dispense.

Pharmacy Information *(Fields marked * are REQUIRED.)*

*Pharmacy Name:

*Address Line #1:

Address Line #2:

*City:

*State:

*ZIP Code:

*National Provider Identifier (NPI):

*Office Phone Number:

Patient Information

*First Name (Legal):

*Last Name (Legal):

*Date of Birth (MM/DD/YYYY):

Prescriber (HCP) Information

*First Name:

*Last Name:

*National Provider Identifier (NPI):

Continued on Next Page

ZENBEXUS™ REMS | Pharmacy Counseling Checklist

For Females Who Can Become Pregnant and Males

Pharmacy Information (cont'd) (Fields marked * are REQUIRED.)

☐ ***I have counseled the female patient who can become pregnant on:**

- ☐ The risk of embryo-fetal toxicity
- ☐ The need to use effective methods of contraception every time they have sex, beginning at least 4 weeks before taking ZENBEXUS™ (iberdromide), while taking ZENBEXUS, and for at least 4 weeks after stopping ZENBEXUS
- ☐ The need to complete pregnancy tests every 4 weeks if regular menstrual cycles or every 2 weeks if irregular menstrual cycles
- ☐ The need to complete a patient survey every 4 weeks
- ☐ The need to immediately contact their prescriber and stop treatment if the patient misses a menstrual period or if pregnancy is suspected
- ☐ Not donating blood during treatment and for 4 weeks following treatment completion with ZENBEXUS
- ☐ Not donating eggs during treatment and for 4 weeks following treatment completion with ZENBEXUS
- ☐ Returning unused ZENBEXUS to a drug take back location
- ☐ Not sharing ZENBEXUS

☐ ***I have counseled the male patient on:**

- ☐ The risk of embryo-fetal toxicity
- ☐ The need to use latex or synthetic male condoms every time they have sex, beginning at least 4 weeks before taking ZENBEXUS, while taking ZENBEXUS, and for at least 4 weeks after stopping ZENBEXUS
- ☐ Not donating sperm during treatment and for 4 weeks following treatment completion with ZENBEXUS
- ☐ Not donating blood during treatment and for 4 weeks following treatment completion with ZENBEXUS
- ☐ The need to complete a patient survey every 4 weeks
- ☐ Returning unused ZENBEXUS to a drug take back location
- ☐ Not sharing ZENBEXUS

*Counselor Name:

*Date: